

420 Rec'd PCT/PTO 1 2 NOV 1999

Multivalent vaccine composition with mixed carrier

The subject of the present invention is a pharmaceutical composition intended for the treatment or prevention of a number of infections caused by pathogenic agents such as bacteria, which comprises, as immunogenic agent, polysaccharides derived from one or more pathogenic agent.

Bacteria as well as fungi such as yeasts incorporate polysaccharides into their surface structure. Thus, the great majority of bacteria are coated with an exudate of a polysaccharide nature which is attached to the bacterium more or less firmly but which is strictly speaking not an envelope. This exudate is called glycocalyx or capsule. Moreover, the outer membrane of Gram-negative bacteria, consists, inter alia, of lipopolysaccharide (LPS). Finally, polysaccharides are also found in the wall of fungi. These polysaccharides are in fact surface antigens which induce an immune response in an infected mammal.

Such polysaccharides are produced on the basis of units in which the constituents and the bonds are defined and which are characteristic of the bacterial or fungal species considered. These repeating units contain the epitopes, that is to say the structures which determine antigenicity.

The polysaccharides of pathogenic micro-organisms are reputed to be good vaccine agents. As they are, they are effective in adults and children over two years. On the other hand, in breast-feeding infants, some are only slightly or not immunogenic and do not induce any immune response. It is possible to overcome this problem by coupling, via covalent bonding, the polysaccharides to a so-called carrier protein such as diphtheria or tetanus toxoid so as to obtain a polysaccharide-carrier protein conjugate.

The same vaccine composition may contain several conjugates. Indeed, the trend is to combine several vaccinal agents intended to prevent or to treat

15 Thus, this is for example the case when it is  
sought to vaccinate against *Streptococcus pneumoniae*  
infections. Pneumococcal infections are a real public  
health problem especially since they are found in the  
severe forms of pneumonia, septicaemia and meningitis.  
20 In industrialized countries, they affect each year 30  
to 100 per 100,000 children under three years. The  
mortality rate in cases of bacteraemia and meningitis  
is 15 to 30% whereas 5% of children die of pneumonia.

35           Moreover, *Streptococcus pneumoniae* is responsible for a number of otitis infections which are more benign but very common. The number of children that have had an otitis infection before the age of six is evaluated at about 75% and the number of otitis

infections caused by pneumococcus at 30 to 50%. In developed countries, otitis infections caused by pneumococcus are due to serogroup 19 in 25% of cases, followed by serogroups/serotypes 23 (13%), 6 and 14 (12%), 9 and 18 (4%) and 4 and 1 (2%).

A pneumococcal vaccine containing the polysaccharides of 23 serotypes is already commercially available. This vaccine makes it possible to effectively combat invasive infections in adults and has a transient action in children over seven months.

The capsular polysaccharides of pneumococci are T-independent antigens, i.e. they can induce antibodies, preferably of the IgM type, without the help of T cells and are not capable of promoting a booster response of the IgG type. When they are coupled to a carrier protein, these polysaccharides then prove capable of inducing a T-dependent response, most particularly in neonates and should provide long-term protection.

Clinical studies have been carried out in Finland and Israel with pneumococcal vaccines having four valencies containing conjugates 6B, 9V, 18C and 23F in which the polysaccharide was coupled either to Dt or to Tt. The doses were 1, 3 or 10 µg of polysaccharide per valency. Each of these formulations was administered simultaneously with an anti-*Haemophilus* vaccine (polyribitolphosphate coupled to Tt; Act-HIB marketed by Pasteur Mérieux Connaught) and an anti-diphtheria, tetanus, whooping cough vaccine (for Finland, D.T.Coq marketed by KTL). Furthermore, these three administrations were carried out accompanied or not by simultaneous administration of an oral or injectable polio vaccine. They were repeated twice at a few weeks interval, and then once, one year after the first immunization.

The results of these studies as reported in the table below have made it possible to demonstrate an effect of negative interference of the diphtheria and

00423698 "021000

5  
10  
15  
20

Israeli study	Anti-HiB antibody in µg/ml
Placebo	6.62
Tetavalent pneumo	
Tt: 3 µg	2.81
Dt: 3 µg	4.62

Accordingly, the subject of the invention is a composition comprising "n" conjugates C1 to Cn, each conjugate being composed (i) of a polysaccharide, in particular a polysaccharide derived from a *Streptococcus pneumoniae* serotype/serogroup S1 to Sn respectively, and (ii) of a carrier protein P1 to Pn respectively, "n" being a number equal to or greater than 2; in which composition the polysaccharides S1 to Sn are identical or different and in which the carrier proteins P1 to Pn are selected independently from a group consisting of "m" carrier proteins A1 to Am, "m" being a number equal to or greater than 2, provided that at least one of the carrier proteins P1 to Pn is different from the others.

According to another aspect, the subject of the invention is also a composition which comprises "n" conjugates C1 to Cn, each conjugate being composed (i) of a polysaccharide S1 to Sn respectively and (ii) a carrier protein P1 to Pn respectively, "n" being a number equal to or greater than 2; in which composition the polysaccharides S1 to Sn are identical or different and in which the carrier proteins P1 to Pn are selected independently from a group consisting of diphtheria (Dt) and tetanus (Tt) toxoids, provided that at least one of the carrier proteins P1 to Pn is different from the others; and which is characterized in that the quantity of Dt and Tt is respectively less than or equal to 200 and 50 µg/dose. In other words, a composition according to the invention comprises one or more polysaccharide conjugates in which the polysaccharide is coupled to the diphtheria toxoid (Dt) and one or more polysaccharide conjugates in which the polysaccharide is coupled to the tetanus toxoid (Tt) and is characterized in that the quantity of Dt and Tt is respectively less than or equal to 200 and 50 µg/dose.

By way of illustration, the following compositions are envisaged:

00423698 "021000





most equal to "n", "n" being as defined above. In particular, persons skilled in the art can determine the minimum number of different carrier proteins which is necessary in order to avoid the phenomenon of interference. To do this, they will take into account the maximum load that should not be exceeded for each of the carrier proteins. "Maximum load" refers to the quantity of carrier protein above which a reduced immune response is observed against one or more polysaccharides compared with a corresponding monovalent composition (conjugate taken separately).

In particular, as regards the diphtheria toxoid and the tetanus toxoid, it is estimated that, advantageously, the quantity of these proteins present in a dose of a composition according to the invention should not exceed 200 and 50 µg respectively, such a dose being envisaged for administration in a mammal, preferably a human. Preferably, the Dt load is less than or equal to 150, 120 or 100 µg, most preferably 80 or 60 µg. Preferably, the Tt load is less than or equal to 35 or 25 µg, most preferably 20 or 10 µg.

Thus, it may be accepted that for a composition using only two different carrier proteins, the selection of these proteins will be made from a group consisting of proteins A1 and A2. Preferably, A1 and A2 may be diphtheria toxoid (Dt) and tetanus toxoid (Tt) respectively or vice versa.

According to a specific embodiment, a composition using only two different carrier proteins is characterized by a balanced distribution of the number of polysaccharides conjugated with the first carrier protein and of the number of polysaccharides conjugated with the second carrier protein. For example, when "n" is an even number, "n"/2 carrier proteins P1 to Pn are A1 and "n"/2 carrier proteins P1 to Pn are A2 or when "n" is an odd number, ("n"+1)/2 carrier proteins P1 to Pn are A1 and ("n"-1)/2 carrier proteins P1 to Pn are A2.

09423698-021000



A polysaccharide useful for the purposes of the present invention may be in particular a polysaccharide of bacterial or fungal origin. It may be in particular a polysaccharide from *Streptococcus* e.g. *Streptococcus pneumoniae*, *Staphylococcus*, *Klebsiella*, *Salmonella* e.g. *Salmonella typhi*, *Escherichia*, *Shigella*, *Neisseria* e.g. *Neisseria meningitidis*, *Haemophilus* e.g. *Haemophilus influenzae*, *Moraxella*, *Vibrio cholerae* or *Mycobacterium tuberculosis*.

In a composition according to the invention, the polysaccharides may be derived from different species or alternatively may all be derived from the same species, e.g. from the same bacterial species, possibly of different serogroups/serotypes. In order to illustrate this last possibility, there may be mentioned a composition according to the invention intended to vaccinate against pneumococcal infections, which contains at least 8 valencies, preferably 10 or 11 valencies chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F.

Thus, a composition constituting a pneumococcal vaccine advantageously comprises 10 or 11 valencies, e.g. represented by 10 or 11 conjugates in which the polysaccharides are all different from each other and are derived (have as origin) serotypes/serogroups chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F of *S. pneumoniae*. It may be in particular a composition which comprises 10 or 11 conjugates selected from:

- serotype 1 polysaccharide coupled to Tt or to Dt;
- serotype 3 polysaccharide coupled to Dt;
- serotype 4 polysaccharide coupled to Tt;
- serotype 5 polysaccharide coupled to Tt or to Dt;
- serotype 6B polysaccharide coupled to Dt;
- serotype 7F polysaccharide coupled to Tt or to Dt;
- serotype 9V polysaccharide coupled to Tt;
- serotype 14 polysaccharide coupled to Dt;
- serotype 18C polysaccharide coupled to Dt;
- serotype 19F polysaccharide coupled to Tt; and

- serotype 23F polysaccharide coupled to Tt.

Under another aspect, a composition constituting a pneumococcal vaccine may comprise 10 or 11 valencies represented by 12 to 22, especially 12 to 15 conjugates, in which the polysaccharides are derived from the serotypes chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F, in which composition conjugates of the same valency differ from each other in the carrier protein. It may be in particular a composition which comprises:

- serotype 1 polysaccharide coupled to Tt;
- serotype 3 polysaccharide coupled to Dt;
- serotype 4 polysaccharide coupled to Tt;
- serotype 5 polysaccharide coupled to Tt;
- serotype 6B polysaccharide coupled to Dt;
- serotype 6B polysaccharide coupled to Tt;
- serotype 7F polysaccharide coupled to Tt;
- serotype 9V polysaccharide coupled to Tt;
- serotype 9V polysaccharide coupled to Dt;
- serotype 14 polysaccharide coupled to Dt;
- serotype 18C polysaccharide coupled to Dt;
- serotype 18C polysaccharide coupled to Tt;
- serotype 19F polysaccharide coupled to Tt;
- serotype 23F polysaccharide coupled to Tt; and
- serotype 23F polysaccharide coupled to Dt.

Such a polysaccharide may be advantageously extracted from the microorganism according to conventional methods and purified likewise. This polysaccharide may be used in the crude form after extraction/purification. Alternatively, it may be fragmented in order to obtain a polysaccharide having an average molecular weight less than that of the polysaccharide originally extracted. A particularly advantageous fermentation method is described in WO 93/7178 which is incorporated by way of reference.

A conjugate in which a polysaccharide is coupled by covalent bonding to a carrier protein may be obtained according to conventional methods well known to persons skilled in the art. It may make use of a

09423698.021000

linker or a spacer to carry out the conjugation. Depending on the mode of conjugation used, the conjugate resulting therefrom may be a conjugate in which the polysaccharide is linked to the protein by a  
5 single chemical function (sun or neoglycoconjugate type), or by several functions (random coil type). It is within the capability of persons skilled in the art to determine the most appropriate mode of conjugation as a function of the nature of the polysaccharide and  
10 more particularly of the chemical groups carried by the polysaccharide which may be used during the conjugation reaction.

A composition according to the invention may be manufactured conventionally. In particular, it may be  
15 formulated with a pharmaceutically acceptable diluent or vehicle, e.g. water or a saline solution. In addition, the composition may contain customary ingredients such as a buffer, a preservative or stabilizer, an adjuvant such as an aluminum compound,  
20 e.g. an aluminium hydroxide, an aluminium phosphate or an aluminium hydroxyphosphate, and, where appropriate, a lyophilization excipient. In general, these products may be selected as a function of the mode and route of administration and based on standard pharmaceutical  
25 practices. Appropriate carriers or diluents as well as what is essential for the preparation of a pharmaceutical composition are described in *Remington's Pharmaceutical Sciences*, a standard reference book in this field.

30 A composition according to the invention may be administered by any conventional route which is used in the field of vaccines, in particular by the systemic, i.e. parenteral, route, e.g. by the subcutaneous, intramuscular, intradermal or intravenous route, or by  
35 the mucosal route, e.g. by the oral or nasal route.

The administration may take place in a single dose or in a dose repeated once or several times, e.g. once, twice or three times, after a certain interval of time. The appropriate dosage will vary as a function of

09423598-021000

various parameters, for example the number of valencies contained in the composition, the nature of the polysaccharide(s) used or the mode of administration. As a guide, it is indicated that good results may be obtained with per valency, a polysaccharide dose of 0.5 to 100 µg, preferably of 1 to 50 µg, most preferably of 1 to 10 µg. A dose of the composition according to the invention may be advantageously in a volume of 0.1 to 2 ml.

There are presented below, by way of example, various pneumococcal vaccines having multiple valencies, the valencies being chosen from serotypes 1, 3, 4, 5, 6B, 7F, 9V, 14, 18C, 19F and 23F. The polysaccharides derived from these serotypes were fragmented according to the method described in WO 93/7178. The polysaccharides coupled to Tt (except the type 1 polysaccharide) were coupled according to the conjugation method described in WO 93/7178. Briefly, a polysaccharide is subjected to reductive amination in the presence of sodium cyanoborohydride in order to link a molecule of diaminoethane to a terminal reducing group. The polysaccharide thus derived is then activated by a succinimide group using disuccinimidyl suberate (DSS). The polysaccharide thus activated is reacted directly with the carrier protein. The serotype 1 polysaccharide was coupled Dt to Tt [sic] according to the conjugation method described in US patent No. 5,204,098 which is incorporated by way of reference. The other polysaccharides coupled to Dt were coupled as follows; hydrazide groups were incorporated onto the polysaccharide by reacting the polysaccharide with an excess of adipic acid dihydrazide (ADH) in the presence of ethyl dimethyl amino propyl carbodiimide (EDAC) and sodium cyanoborohydride (for all the types except 3) or simply in the presence of sodium cyanoborohydride (for type 3). The polysaccharide thus derived is reacted with the carrier protein in the presence of EDAC. The experimental conditions were controlled so as to obtain conjugates in which the quantity of protein is between

09423698-021000

one and four times, preferably twice, the value of the quantity of polysaccharide. Thus, for a dose, 3 µg of a particular polysaccharide are coupled to about 6 µg of Dt and 1 µg of a particular polysaccharide is coupled to about 2 µg of Tt.

The Dt and Tt used were prepared by detoxification with formaldehyde starting with toxins extracted respectively from *Corynebacterium diphtheriae* and *Clostridium tetani*.

The formulations contain a phosphate buffer (0.475 mg of  $\text{PO}_4^{2-}$  ion per dose) and sodium chloride (4.5 mg per dose) and may be supplemented with aluminium hydroxide (alum) adjuvant (300 µg of  $\text{Al}^{3+}$  ion per dose) and contain a preservative such as phenoxyethanol formalin. A dose is in the volume of 0.5 ml.

EXAMPLE 1: Octavalent formulation

Carrier protein	Polysaccharide	
	Serotype	Quantity per single dose
Dt	3	3 µg
Tt	4	1 µg
Dt	6B	10 µg
Tt	9V	1 µg
Dt	14	3 µg
Dt	18C	3 µg
Tt	19F	1 µg
Tt	23F	1 µg

EXAMPLE 2: Formulations F3, F4 and F3bis containing 11 valencies

Carrier Protein	Polysaccharide			
	Serogroup/ Serotype	Qty per single dose of formu- lation F3 (µg)	Qty per single dose of formu- lation F4 (µg)	Qty per single dose of formu- lation F3 bis (µg)
Tt	1	1	-	1
Dt	1	-	3	-
Dt	3	3	3	3
Tt	4	1	1	1
Tt	5	1	-	1
Dt	5	-	3	
Dt	6B	10	10	3
Tt	6B	-	-	1
Tt	7F	1	-	1
Dt	7F	-	3	-
Tt	9V	1	1	1
Dt	9V	-	-	3
Dt	14	3	3	3
Dt	18C	3	3	3
Tt	18C	-	-	1
Tt	19F	1	1	1
Tt	23F	1	1	1
Dt	23F	-	-	3

- 5                      The approximate protein load in each of the three formulations is as follows:

	F3	F4	F3 bis
Dt	about 40 µg	about 60 µg	about 40 µg
Tt	about 15 µg	about 8 µg	about 18 µg

- 10                      (which corresponds to a protein/polysaccharide weight ratio of about 2).